

THE 1, 2, 3-CONJECTURE AND 1, 2-CONJECTURE FOR SPARSE GRAPHS

DANIEL W. CRANSTON

Virginia Commonwealth University
Richmond, VA, USA

e-mail: dcranston@vcu.edu

SOGOL JAHANBEKAM¹

University of Colorado Denver
Denver, CO, USA

e-mail: sogol.jahanbekam@ucdenver.edu

AND

DOUGLAS B. WEST²

Zhejiang Normal University, Jinhua, China
and University of Illinois, Urbana, IL, USA

e-mail: west@math.uiuc.edu

Abstract

The 1, 2, 3-Conjecture states that the edges of a graph without isolated edges can be labeled from $\{1, 2, 3\}$ so that the sums of labels at adjacent vertices are distinct. The 1, 2-Conjecture states that if vertices also receive labels and the vertex label is added to the sum of its incident edge labels, then adjacent vertices can be distinguished using only $\{1, 2\}$. We show that various configurations cannot occur in minimal counterexamples to these conjectures. Discharging then confirms the conjectures for graphs with maximum average degree less than $8/3$. The conjectures are already confirmed for larger families, but the structure theorems and reducibility results are of independent interest.

Keywords: 1, 2, 3-Conjecture, 1, 2-Conjecture, reducible configuration, discharging method.

2010 Mathematics Subject Classification: 05C15, 05C22, 05C78.

¹Research supported in part by National Science Foundation grant DMS 09-01276.

²Research supported in part by National Security Agency grant H98230-10-1-0363.

REFERENCES

- [1] L. Addario-Berry, K. Dalal, C. McDiarmid, B.A. Reed and A. Thomason, *Vertex-colouring edge-weightings*, *Combinatorica* **27** (2007) 1–12.
doi:10.1007/s00493-007-0041-6
- [2] L. Addario-Berry, K. Dalal and B.A. Reed, *Degree constrained subgraphs*, *Discrete Appl. Math.* **156** (2008) 1168–1174.
doi:10.1016/j.dam.2007.05.059
- [3] N. Alon, *Combinatorial Nullstellensatz*, *Combin. Probab. Comput.* **8** (1999) 7–29.
doi:10.1017/S0963548398003411
- [4] T. Bartnicki, J. Grytczuk and S. Niwczyk, *Weight choosability of graphs*, *J. Graph Theory* **60** (2009) 242–256.
doi:10.1002/jgt.20354
- [5] M. Kalkowski, *A note on the 1, 2-conjecture*, submitted (also in Ph.D. Thesis, 2009).
- [6] M. Kalkowski, M. Karoński and F. Pfender, *Vertex-coloring edge-weightings: towards the 1-2-3-conjecture*, *J. Combin. Theory (B)* **100** (2010) 347–349.
doi:10.1016/j.jctb.2009.06.002
- [7] M. Karoński, T. Łuczak and A. Thomason, *Edge weights and vertex colours*, *J. Combin. Theory (B)* **91** (2004) 151–157.
doi:10.1016/j.jctb.2003.12.001
- [8] J. Przybyło and M. Woźniak, *On a 1, 2 conjecture*, *Discrete Math. Theoret. Comput. Sci.* **12** (2010) 101–108.
- [9] B. Seamone, *The 1-2-3 conjecture and related problems: a survey*, submitted (<http://arxiv.org/abs/1211.5122>).
- [10] T. Wang and Q. Yu, *On vertex-coloring 13-edge-weighting*, *Front. Math. China* **3** (2008) 581–587.
doi:10.1007/s11464-008-0041-x
- [11] T.-L. Wong, X. Zhu and D. Yang, *List total weighting of graphs*, in: G.O.H. Katona, A. Schrijver, T. Szőnyi and G. Sági, Eds., *Fete of Combinatorics and Computer Science*, Bolyai Soc. Math. Stud., vol. 20 (Springer, Berlin, Heidelberg, 2010) 337–353.
doi:10.1007/978-3-642-13580-4_13
- [12] T.-L. Wong and X. Zhu, *Total weight choosability of graphs*, *J. Graph Theory* **66** (2011) 198–212.
doi:10.1002/jgt.20500
- [13] T.-L. Wong and X. Zhu, *Every graph is (2, 3)-choosable*, submitted.

Received 29 March 2013

Revised 13 November 2013

Accepted 13 November 2013