

ON THE UNRECOGNIZABILITY BY PRIME GRAPH FOR THE ALMOST SIMPLE GROUP $\text{PGL}(2, 9)$

ALI MAHMOUDIFAR

*Department of Mathematics, Tehran North Branch
Islamic Azad University, Tehran, Iran*

e-mail: alimahmoudifar@gmail.com

Abstract

The prime graph of a finite group G is denoted by $\Gamma(G)$. Also G is called recognizable by prime graph if and only if each finite group H with $\Gamma(H) = \Gamma(G)$, is isomorphic to G . In this paper, we classify all finite groups with the same prime graph as $\text{PGL}(2, 9)$. In particular, we present some solvable groups with the same prime graph as $\text{PGL}(2, 9)$.

Keywords: projective general linear group, prime graph, recognition.

2010 Mathematics Subject Classification: 20D05, 20D60.

1. INTRODUCTION

Let n be a natural number. We denote by $\pi(n)$, the set of all prime divisors of n . Also Let G be a finite group. The set $\pi(|G|)$ is denoted by $\pi(G)$. The set of element orders of G is denoted by $\pi_e(G)$. We denote by $\mu(S)$, the maximal numbers of $\pi_e(G)$ under the divisibility relation. The *prime graph* of G is a graph whose vertex set is $\pi(G)$ and two distinct primes p and q are joined by an edge (and we write $p \sim q$), whenever G contains an element of order pq . The prime graph of G is denoted by $\Gamma(G)$. A finite group G is called *recognizable by prime graph* if for every finite group H such that $\Gamma(G) = \Gamma(H)$, then we have $G \cong H$. So G is *recognizable by prime graph* whenever there exists a fin finite group K such that $\Gamma(K) = \Gamma(G)$ in while K is not isomorphic to G .

For the almost simple group $\text{PGL}(2, q)$, there are a lot of results about the recognition by prime graph. In [8], it is proved that if p is a prime number which is not a Mersenne or Fermat prime and $p \neq 11, 19$ and $\Gamma(G) = \Gamma(\text{PGL}(2, p))$, then G has a unique nonabelian composition factor which is isomorphic to $\text{PSL}(2, p)$ and if $p = 13$, then G has a unique nonabelian composition factor which is

isomorphic to $\text{PSL}(2, 13)$ or $\text{PSL}(2, 27)$. We know that $\text{PGL}(2, 2^\alpha) \cong \text{PSL}(2, 2^\alpha)$. For the characterization of such simple groups we refer to [9, 10]. In [1], it is proved that if $q = p^\alpha$, where p is an odd prime and α is an odd natural number, then $\text{PGL}(2, q)$ is uniquely determined by its prime graph.

By the above description, we get that the characterization by prime graph of $\text{PGL}(2, p^k)$, where p is an odd prime number and k is even, is an open problem. In this paper as the main result we consider the recognition by prime graph of the almost simple groups $\text{PGL}(2, 3^2)$. Moreover, we construct some solvable group with the same prime graph as $\text{PGL}(2, 3^2)$.

2. PRELIMINARY RESULTS

Lemma 2.1. *Let G be a finite group and $N \trianglelefteq G$ such that G/N is a Frobenius group with kernel F and cyclic complement C . If $(|F|, |N|) = 1$ and F is not contained in $NC_G(N)/N$, then $p|C| \in \pi_e(G)$ for some prime divisor p of $|N|$.*

Lemma 2.2. *Let G be a Frobenius group with kernel F and complement C . Then the following assertions hold:*

- (a) F is a nilpotent group.
- (b) $|F| \equiv 1 \pmod{|C|}$.
- (c) Every subgroup of C of order pq , with p, q (not necessarily distinct) primes, is cyclic.

In particular, every Sylow subgroup of C of odd order is cyclic and a Sylow 2-subgroup of C is either cyclic or generalized quaternion group. If C is a non-solvable group, then C has a subgroup of index at most 2 isomorphic to $SL(2, 5) \times M$, where M has cyclic Sylow p -subgroups and $(|M|, 30) = 1$.

By using [13, Theorem A] we have the following result:

Lemma 2.3. *Let G be a finite group with $t(G) \geq 2$. Then one of the following holds:*

- (a) G is a Frobenius or 2-Frobenius group;
- (b) there exists a nonabelian simple group S such that $S \leq \overline{G} := G/N \leq \text{Aut}(S)$ for some nilpotent normal π_1 -subgroup N of G and \overline{G}/S is a π_1 -group.

3. MAIN RESULTS

Lemma 3.1. *There exists a Frobenius group $G = K : C$, where K is an abelian 3-group and $\pi(C) = \{2, 5\}$, such that $\Gamma(G) = \Gamma(\text{PGL}(2, 9))$.*

Proof. Let F be a finite field with 3^4 elements. Also let V be the additive group of F and H be the multiplicative group $F \setminus \{0\}$. We know that H acts on V by right product. So $G := V \rtimes H$ is a finite group such that $\pi(G) = \{2, 3, 5\}$, since $|V| = 3^4$ and $|H| = 80$. On the other hand H acts fixed point freely on V , so G is a Frobenius group with kernel V and complement H . Since the multiplicative group $F \setminus \{0\}$ is cyclic, H is cyclic too. Therefore, G has an element of order 10, and so the prime graph of G consists just one edge, which is the edge between 2 and 5. This implies that $\Gamma(G) = \Gamma(\text{PGL}(2, 9))$, as desired. ■

Lemma 3.2. *There exists a Frobenius group $G = K : C$, where $\pi(K) = \{2, 5\}$ and C is a cyclic 3-group such that $\Gamma(G) = \Gamma(\text{PGL}(2, 9))$.*

Proof. Let F_1 and F_2 be two fields with 2^2 and 5^2 elements, respectively. Let $V := F_1 \times F_2$ be the direct product of the additive groups F_1 and F_2 . Also let $H := H_1 \times H_2$, be the direct product of H_1 and H_2 , which are the multiplicative groups $F_1 \setminus \{0\}$ and $F_2 \setminus \{0\}$, respectively. We know that H_i acts fixed point freely on F_i , where $1 \leq i \leq 2$. So we define an action of H on V as follows: for each $(h, h') \in H$ and $(g, g') \in V$, we define $(g, g')^{(h, h')} := (hg, h'g')$. It is clear that this definition is well defined. So we may construct a finite group $G = V \rtimes H$. On the other hand H acts fixed point freely on V . So G is a Frobenius group with kernel V and complement H such that $\pi(V) = \{2, 5\}$ and $\pi(H) = \{3\}$. Finally, since V is nilpotent, we get that $\Gamma(G)$ contains an edge between 2 and 5, so $\Gamma(G) = \Gamma(\text{PGL}(2, 9))$. ■

Lemma 3.3. *There exists a 2-Frobenius group G with normal series $1 \triangleleft H \triangleleft K \triangleleft G$, such that $\pi(H) = \{5\}$, $\pi(G/K) = \{2\}$ and $\pi(K/H) = \{3\}$, such that $\Gamma(G) = \Gamma(\text{PGL}(2, 9))$.*

Proof. Let F be a field with 5^2 elements and V be its additive group. We know that $F = \{0, 1, \alpha, \alpha^2, \dots, \alpha^{23}\}$, where α is a generator of the multiplicative group $F \setminus \{0\}$. Hence $|\alpha| = 24$, and so $\beta := \alpha^8$ has order 3. Also $\langle \beta \rangle \cong Z_3$ and $\text{Aut}(Z_3) \cong Z_2$. This argument implies that we may construct a Frobenius group $T := \langle \beta \rangle : \langle \gamma \rangle$, where γ is an involution.

Now we define an action of T on V as follows: for each $\beta^x \gamma^y \in T$ and $v \in V$, $v^{\beta^x \gamma^y} := \beta^x v$, where $1 \leq x \leq 3$ and $1 \leq y \leq 2$. Therefore $G := V : T$ is a 2-Frobenius group with desired properties. ■

Theorem 3.4. *Let G be a finite group. Then $\Gamma(G) = \Gamma(\text{PGL}(2, 3^2))$ if and only if G is isomorphic to one of the following groups:*

- (1) A Frobenius group $K : C$, where K is an abelian 3-group and $\pi(C) = \{2, 5\}$,
- (2) A Frobenius group $K : C$, where $\pi(K) = \{2, 5\}$ and C is a cyclic 3-group,

- (3) A 2-Frobenius group with normal series $1 \triangleleft H \triangleleft K \triangleleft G$, such that $\pi(H) \subseteq \{2, 5\}$, $\pi(G/K) = \{2\}$ and $\pi(K/H) = \{3\}$,
- (4) Almost simple group $\text{PGL}(2, 3^2)$.

Proof. Throughout the proof, we assume that G is a finite group with the same prime graph as the almost simple group $\text{PGL}(2, 3^2)$. First we note that by [16, Lemma 7], we have:

$$\mu(\text{PGL}(2, 9)) = \{3, 8, 10\}.$$

Hence in $\Gamma(\text{PGL}(2, 3^2))$ (and so in $\Gamma(G)$), there is only one edge which is the edge between 2 and 5 and so 3 is an isolated vertex. This implies that $\Gamma(G)$ has two connected components $\{3\}$ and $\{2, 5\}$. Thus by Lemma 2.3, we get that G is a Frobenius group or 2-Frobenius group or there exists a nonabelian simple group S such that $S \leq G/\text{Fit}(G) \leq \text{Aut}(S)$. We consider each possibility for G .

Let G be a Frobenius group with kernel K and complement C . We know that K is nilpotent and C is a connected component of the prime graph of G . Also by the above description, 3 is not adjacent to 2 and 5 in $\Gamma(G)$. This shows that either $\pi(K) = \{3\}$ or $\pi(C) = \{3\}$. We consider these cases, separately.

Case 1. Let $\pi(K) = \{3\}$. Hence the order of complement C , is even and so K is an abelian subgroup of G . Also by the above description, $\pi(C) = \{2, 5\}$. Since C is a connected component of $\Gamma(G)$, there is an edge between 2 and 5. So it follows that $\Gamma(G) = \Gamma(\text{PGL}(2, 9))$, which implies groups satisfying in (1).

Case 2. Let $\pi(C) = \{3\}$. Hence $\pi(K) = \{2, 5\}$. Since K is nilpotent, we get that 2 and 5 are adjacent in $\Gamma(G)$. This means $\Gamma(G) = \Gamma(\text{PGL}(2, 9))$ and so we get (2).

Case 3. Let G be a 2-Frobenius group with normal series $1 \triangleleft H \triangleleft K \triangleleft G$. Since $\pi(K/H)$ and $\pi(H) \cup \pi(G/K)$ are the connected components of $\Gamma(G)$, we get that $\pi(K/H) = \{3\}$ and $\pi(H) \cup \pi(G/K) = \{2, 5\}$. This implies (3).

Case 4. Let there exist a nonabelian simple group S , such that $S \leq \bar{G} := G/\text{Fit}(G) \leq \text{Aut}(S)$. Since $\pi(S) \subseteq \pi(G)$, $\pi(S) = \{2, 3, 5\}$. The finite simple groups with this property are classified in [12, Table 8]. So we get that S is isomorphic to one of the simple groups A_5 , $\text{PSU}(4, 2)$ and $\text{PSL}(2, 9) (\cong A_6)$.

Subcase 4.1. Let $S \cong A_5$. We know that $\text{Aut}(A_5) = S_5$. So \bar{G} is isomorphic to the alternating group A_5 or the symmetric group S_5 . Since in the prime graph of S_5 , there is an edge between 2 and 3, hence we get that \bar{G} is not isomorphic to S_5 . Thus, $G/\text{Fit}(G) = A_5$. In the prime graph of A_5 , 2 and 5 are nonadjacent. Then at least one of the prime numbers 2 or 5, belongs to $\pi(\text{Fit}(G))$.

Let $5 \in \pi(\text{Fit}(G))$. Let F_5 be a Sylow 5-subgroup of $\text{Fit}(G)$. Since F_5 is a characteristic subgroup of $\text{Fit}(G)$ and $\text{Fit}(G)$ is a normal subgroup of G , $F_5 \trianglelefteq G$.

On the other hand in alternating group A_5 , the subgroup $\langle (12)(34), (13)(24) \rangle : \langle (123) \rangle$ is a Frobenius subgroup isomorphic to $2^2 : 3$. We recall that by the previous argument, $F_5 \trianglelefteq G$, and so G has a subgroup isomorphic to $5^\alpha : (2^2 : 3)$. So by Lemma 2.1, we get that 3 is adjacent to 5, a contradiction.

Subcase 4.2. Let $S \cong \text{PSU}(4, 2)$. By [4], there is an edge between 3 and 2 which is a contradiction.

Subcase 4.3. Let $S \cong \text{PSL}(2, 9)$. Then \bar{G} is isomorphic to $\text{PSL}(2, 9)$ or $\text{PSL}(2, 9) : \langle \theta \rangle$, where θ is a diagonal, field or diagonal-field automorphism of $\text{PSL}(2, 9)$. In particular θ is an involution. If θ is a field or diagonal-field automorphism of $\text{PSL}(2, 9)$, then the semidirect product $\text{PSL}(2, 9) : \langle \theta \rangle$ contains an element of order 6. Hence θ is neither a field automorphism nor a diagonal-field automorphism. Therefore θ is a diagonal automorphism and so $\bar{G} \cong \text{PGL}(2, 9)$.

By the above discussion, $G/\text{Fit}(G) \cong \text{PGL}(2, 9)$. It is enough to prove that $\text{Fit}(G) = 1$. On the contrary, let $r \in \pi(\text{Fit}(G))$. Also let F_r be the Sylow r -subgroup of $\text{Fit}(G)$. Since $\text{Fit}(G)$ is nilpotent, we can write $\text{Fit}(G) = O_{r'}(\text{Fit}(G)) \times F_r$. So if we put $\tilde{G} = G/O_{r'}(\text{Fit}(G))$, then we get that:

$$\text{PGL}(2, 9) \cong \frac{G}{\text{Fit}(G)} \cong \frac{\tilde{G}}{F_r} \cong \frac{\tilde{G}/\Phi(F_r)}{F_r/\Phi(F_r)}.$$

Since $F_r/\Phi(F_r)$ is an elementary abelian group, without loose of generality we may assume that $F := \text{Fit}(G)$ is an elementary abelian r -group and $G/F \cong \text{PGL}(2, 9)$.

If $r = 3$, then by , we conclude that in $\Gamma(G)$, 2 and 3 are adjacent, which is a contradiction. So let $r \neq 3$. Also let S_3 be a Sylow 3-subgroup of $\text{PGL}(2, 9)$. We know that S_3 is not cyclic. On the other hand $F \rtimes S_3$ is a Frobenius group since 3 is an isolated vertex of $\Gamma(G)$. This follows that S_3 is cyclic which is impossible. Therefore $F = 1$ and so $G \cong \text{PGL}(2, 9)$, which completes the proof. ■

REFERENCES

- [1] Z. Akhlaghi, M. Khatami and B. Khosravi, *Characterization by prime graph of $\text{PGL}(2, p^k)$ where p and k are odd*, Int. J. Alg. Comp. **20** (7) (2010) 847–873.
doi:10.1142/S021819671000587X
- [2] S. Shariati Beynekalae, A. Iranmanesh and M. Foroudi Ghasemabadi, *Quasirecognition by prime graph of the simple group $B_n(2)$* , Southeast Asian Bull. Math. **39** (2015) 181–193.
- [3] L. Wang, *Quasirecognition of $L_{22}(2)$ by Its prime graph*, Southeast Asian Bull. Math. **35** (2015) 883–889.
- [4] A.A. Buturlakin, *Spectra of finite linear and unitary groups*, Algebra and Logic **47** (2) (2008) 91–99.
doi:10.1007/s10469-008-9003-3

- [5] J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker and R.A. Wilson, *Atlas of Finite Groups* (Oxford University Press, Oxford, 1985).
- [6] K.W. Gruenberg and K.W. Roggenkamp, *Decomposition of the augmentation ideal and of the relation modules of a finite group*, Proc. London Math. Soc. **31** (2) (1975) 149–166.
doi:10.1112/plms/s3-31.2.149
- [7] M. Hagie, *The prime graph of a sporadic simple group*, Comm. Algebra **31** (9) (2003) 4405–4424. doi:10.1081/AGB-120022800
- [8] M. Khatami, B. Khosravi and Z. Akhlaghi, *NCF-distinguishability by prime graph of $PGL(2, p)$, where p is a prime*, Rocky Mountain J. Math. **4** (5) (2011) 1523–1545.
doi:10.1216/rmj-2011-41-5-1523
- [9] B. Khosravi, *n -Recognition by prime graph of the simple group $PSL(2, q)$* , J. Algebra Appl. **7** (6) (2008) 735–748.
doi:10.1142/S0219498808003090
- [10] B. Khosravi, B. Khosravi and B. Khosravi, *2-Recognizability of $PSL(2, p^2)$ by the prime graph*, Siberian Math. J. **49** (4) (2008) 749–757.
doi:10.1007/s11202-008-0072-2
- [11] B. Khosravi, B. Khosravi and B. Khosravi, *On the prime graph of $PSL(2, p)$ where $p > 3$ is a prime number*, Acta. Math. Hungarica **116** (4) (2007) 295–307.
doi:10.1007/s10474-007-6021-x
- [12] R. Kogani-Moghadam and A.R. Moghaddamfar, *Groups with the same order and degree pattern*, Sci. China Math. **55** (4) (2012) 701–720.
doi:10.1007/s11425-011-4314-6
- [13] A.S. Kondrat'ev, *Prime graph components of finite simple groups*, Math. USSR-SB. **67** (1) (1990) 235–247.
doi:10.1070/SM1990v067n01ABEH001363
- [14] V.D. Mazurov, *Characterizations of groups by arithmetic properties*, Proceedings of the International Conference on Algebra, Algebra Colloq. **11** (1) (2004) 129–140.
- [15] V.D. Mazurov, *Characterizations of finite groups by sets of their element orders*, Algebra Logic **36** (1) (1997) 23–32.
doi:10.1007/BF02671951
- [16] A.R. Moghaddamfar and W.J. Shi, *The number of finite groups whose element orders is given*, Beitrage Algebra Geom. **47** (2) (2006) 463–479.
- [17] J.S. Williams, *Prime graph components of finite groups*, J. Algebra **69** (2) (1981) 487–513. doi:10.1016/0021-8693(81)90218-0
- [18] A.V. Zavarnitsin, *Recognition of finite groups by the prime graph*, Algebra Logic **43** (4) (2006) 220–231. doi:10.1007/s10469-006-0020-9
- [19] K. Zsigmondy, *Zur theorie der potenzreste*, Monatsh. Math. Phys. **3** (1892) 265–284.
doi:10.1007/BF01692444

Received 9 September 2016

Revised 20 October 2016