

THE RINGS WHICH ARE BOOLEAN*

IVAN CHAJDA

Department of Algebra and Geometry
Palacký University Olomouc, 17. listopadu 12
771 46 Olomouc, Czech Republic

e-mail: ivan.chajda@upol.cz

AND

FILIP ŠVRČEK

Department of Algebra and Geometry
Palacký University Olomouc, 17. listopadu 12
771 46 Olomouc, Czech Republic

e-mail: filip.svrcek@upol.cz

Abstract

We study unitary rings of characteristic 2 satisfying identity $x^p = x$ for some natural number p . We characterize several infinite families of these rings which are Boolean, i.e., every element is idempotent. For example, it is in the case if $p = 2^n - 2$ or $p = 2^n - 5$ or $p = 2^n + 1$ for a suitable natural number n . Some other (more general) cases are solved for p expressed in the form $2^q + 2m + 1$ or $2^q + 2m$ where q is a natural number and $m \in \{1, 2, \dots, 2^q - 1\}$.

Keywords: Boolean ring, unitary ring, characteristic 2.

2010 Mathematics Subject Classification: 06E20, 16R40.

REFERENCES

- [1] I.T. Adamson, Rings, modules and algebras (Oliver&Boyd, Edinburgh, 1971).
- [2] G. Birkhoff, Lattice Theory, 3rd edition (AMS Colloq. Publ. 25, Providence, RI, 1979).

*This work is supported by the Research and Development Council of the Czech Government via the project MSM6198959214.

- [3] I. Chajda and F. Švrček, *Lattice-like structures derived from rings*, Contributions to General Algebra, Proc. of Salzburg Conference (AAA81), J. Hayn, Klagenfurt **20** (2011), 11–18.
- [4] N. Jacobson, *Structure of Rings* (Amer. Math. Soc., Colloq. Publ. 36 (rev. ed.), Providence, RI, 1964).
- [5] J. Lambek, *Lectures on Rings and Modules* (Blaisdell Publ. Comp., Waltham, Massachusetts, Toronto, London, 1966).
- [6] N.H. McCoy, *Theory of Rings* (Mainillan Comp., New York, 1964).

Received 17 February 2011

Revised 22 June 2011