

ESTIMATES FOR THE DISTRIBUTION OF THE FIRST EXIT TIME OF α -STABLE PROCESSES

WIESŁAW CUPAŁA

Institute of Mathematics
University of Zielona Góra
Podgórna, 50 65–246 Zielona Góra, Poland
e-mail: w.cupala@im.uz.zgora.pl

Abstract

The Varopoulos-Hardy-Littlewood theory and the spectral analysis are used to estimate the tail of the distribution of the first exit time of α -stable processes.

Keywords: first exit time, α -stable processes.

2000 Mathematics Subject Classification: 60G40, 60J25.

1. PRELIMINARIES

Let $\xi(t)$ ($t > 0$) be an α -stable process on \mathbf{R}^d ($d \geq 3$) i.e., it is a Markov process with the strong Markov property, the transition kernel of which is given by the convolution with a function $p_t(x)$ and the Fourier transform of $p_t(x)$ has a form

$$\hat{p}_t(y) = \exp(-t\|y\|^\alpha), \quad (1 \leq \alpha \leq 2).$$

The uniformity in time of α -stable processes implies that the family of operators

$$T_t f(x) = E f(\xi_x(t)), \quad (f \in L^p(\mathbf{R}^d))$$

forms a semigroup of contractions.

Let $f \in L^p(\mathbf{R}^d)$ ($1 \leq p \leq \infty$) and $\frac{1}{p} + \frac{1}{q} = 1$. The form of \hat{p}_t gives us the following estimation:

$$\|T_t f\|_\infty \leq \|p_t\|_q \|f\|_p.$$

Because

$$p_t(x) = \frac{1}{t^{d/\alpha}} p_1\left(\frac{x}{t^{1/\alpha}}\right),$$

we obtain that

$$\|p_t\|_q = t^{-\frac{dp}{\alpha}} \|p_1\|_q,$$

for any $1 \leq q \leq +\infty$. This implies

$$(1) \quad \|T_t f\|_\infty \leq C t^{-\frac{dp}{\alpha}} \|f\|_p \quad (p \geq 1),$$

where the constant C depends only on p .

Let U be a bounded domain in \mathbf{R}^d . Let ξ starts in $x \in U$ (let us denote this fact by $\xi_x(t)$). Trajectories of such processes are right-continuous and have left-hand side limits, so we can define the first exit time from U by

$$\tau_x = \inf\{t \mid \xi_x(t) \notin U\}.$$

Lemma 1. *Let U be a domain in \mathbf{R}^d and $\xi_x(t)$ be an α -stable process which starts in $x \in U$. Let us denote by*

$$I_x(t) = \mathbf{1}_{\{t < \tau_x\}}$$

the indicator of the set of all trajectories for which $t < \tau_x$. Let $f \in L^p(U)$, $1 \leq p \leq \infty$. The family of operators defined by

$$(2) \quad S_t f(x) = E f(\xi_x(t)) I_x(t)$$

forms a semigroup of contractions on $L^p(U)$.

Proof. By using of the strong Markov property of $\xi(t)$ we have

$$I_x(t)I_{\xi_x(t)}(s) = I_x(t+s).$$

Hence

$$\begin{aligned} S_t \circ S_s f(x) &= E S_s f(\xi_x(t)) I_x(t) = \\ &= E(E f(\xi_{\xi_x(t)}(s)) I_{\xi_x(t)}(s)) I_x(t) = \\ &= E f(\xi_x(t)) I_x(t+s). \end{aligned}$$

So, $S_t \circ S_s = S_{t+s}$. ■

The estimation (1) implies

Lemma 2. *Let $f \in L^p(U)$, $1 \leq p \leq \infty$. Then*

$$(3) \quad \|S_t f\|_\infty \leq C t^{\frac{-dp}{\alpha}} \|f\|_p,$$

where constant C depends only on p, d, α .

Let D be the infinitesimal generator of S_t and let $N(\lambda)$ denote the dimension of its spectral projector $P(-\infty, \lambda)$.

Lemma 3. *For any bounded domain U there exists a constant C such that for every $\lambda \geq 0$*

$$N(\lambda) \leq C \lambda^{\frac{d}{\alpha}}.$$

Proof. By (3) and the Varopoulos theory ([2], Theorem 1) we have

$$\|f\|_{2d/(d-\alpha)} \leq C \|D^{\frac{1}{2}} f\|_2, \quad f \in \text{Dom} \left(D^{\frac{1}{2}} \right).$$

So, by the Levin–Solomyak generalization of the CLR inequality (see [1]),

for any $V \geq 0$, $V \in L^{\frac{d}{\alpha}}(U)$ the number of the negative eigenvalues of the operator $D - V$ has an upper bound equals to

$$C_1 \int_U V^{\frac{d}{\alpha}} dx,$$

with some constant C_1 which depends only on α and U . Hence, for any $\lambda > 0$, the operator $D - \lambda Id$ has a finite number of the negative eigenvalues and, because this number is equal to $N(\lambda)$, we have

$$N(\lambda) \leq C_1 |U| \lambda^{\frac{d}{\alpha}}.$$

■

Let (X, μ) be a σ -finite measure space and let Q_t ($t > 0$) be a submarkovian (strongly continuous) symmetric semigroup, i.e., for all $t > 0$, $Q_t : L^2(X) \rightarrow L^2(X)$ is a symmetric operator, and for all $f \in L^2$ with $0 \leq f \leq 1$ we have $0 \leq Q_t f \leq 1$ (we note that S_t is such a semigroup).

Definition 1. Let $u(t, x)$ ($x \in X$, $t > 0$) be a function on $(0, +\infty) \times X$. Let $u(t, \cdot) \in L^1 + L^\infty(X)$. We say that u is a subharmonic function (with respect to the semigroup Q_t) if

$$Q_t u(s, \cdot) \geq u(t + s, \cdot), \quad t, s > 0$$

The proof of the following lemma can be found in [2] (Theorem 2).

Lemma 4. Let $(Q_t; t > 0)$ be a submarkovian symmetric semigroup and let $C, n > 0$ and $1 \leq p < +\infty$ be such that

$$\|Q_t f\|_\infty \leq C t^{-n/2p} \|f\|_p; \quad t > 0, \quad f \in L^p.$$

Then for every subharmonic function $u(t, x)$ and every $0 < s < r \leq +\infty$ we have

$$(4) \quad t^{n/2s} \|u(t, \cdot)\|_r \leq c t^{n/2r} \sup_t \|u(t, \cdot)\|_s,$$

where constant c depends only on C, n, p, r, s .

2. THE MAIN RESULT

Lemma 3 implies that there exists a sequence of positive numbers $w_1 \leq w_2 \leq \dots$ and an orthonormal basis ϕ_1, ϕ_2, \dots in $L^2(U)$ for which

$$D\phi_n = w_n\phi_n.$$

Lemma 5. *There exists $C > 0$ such that*

$$\sup_x |\phi_n| \leq Cw_n^{\frac{2d}{\alpha}}.$$

Proof. Let us notice that

$$S_t\phi_n = \exp(-tw_n)\phi_n.$$

So,

$$u_n(x, t) = \exp(-tw_n)\phi_n(x)$$

is a harmonic function. Lemmas 2 and 4 give us that

$$\sup_x |\exp(-tw_n)\phi_n(x)| \leq Ct^{\frac{-2d}{\alpha}} \sup_t \left(\int_U |\exp(-tw_n)\phi_n(x)|^2 dx \right)^{1/2}.$$

Thus

$$\sup_x |\phi_n(x)| \leq C \inf_t \left(t^{\frac{-2d}{\alpha}} \exp(tw_n) \right).$$

Hence

$$\|\phi_n\|_\infty \leq Cw_n^{\frac{2d}{\alpha}}.$$

■

Theorem 1. *Let τ_x be a first exit time from a bounded domain U . Let w_1 be the smallest eigenvalue of the infinitesimal generator of S_t . Then for every $\varepsilon > 0$ there exists $C(\varepsilon) > 0$ such that for every $t > \varepsilon$*

$$P(t < \tau_x) \leq C(\varepsilon) \exp(-w_1 t).$$

Proof. Let us notice that

$$P(t < \tau_x) \leq \|S_t \mathbf{1}_U\|_\infty,$$

and that

$$\mathbf{1}_U = \sum_{n=1}^{\infty} a_n \phi_n,$$

where

$$\sum_{n=1}^{\infty} |a_n|^2 = \|\mathbf{1}_U\|_2^2 = |U|,$$

and the series is convergent in L^2 . So,

$$(5) \quad S_t \mathbf{1}_U = \sum_{n=1}^{\infty} a_n \exp(-tw_n) \phi_n.$$

By using Lemma 5 we have that (5) converges uniformly on U and

$$\|S_t \mathbf{1}_U\|_\infty \leq C \sum_{n=1}^{\infty} \exp(-w_n t) w_n^{\frac{2d}{\alpha}}.$$

Since,

$$\sum_{n=1}^{\infty} \exp(-w_n t) w_n^{\frac{2d}{\alpha}} \leq C(\varepsilon) \exp(-tw_1)$$

we ended the proof. ■

Corollary 1. *For every $\varepsilon > 0$*

$$E \exp((w_1 - \varepsilon)\tau_x) < +\infty.$$

REFERENCES

- [1] D. Levin and M. Solomyak, *Rozenblum-Lieb-Cwikel inequality for Markov generators*, J. d'Anal. Math. **71** (1997), 173–193.
- [2] N.T. Varopoulos, *Hardy-Littlewood theory for semigroups*, J. Funct. Anal. **63** (1985), 240–260.

Received 25 October 2002