

## SELECTIVE GENERALIZED $F$ TESTS

CÉLIA NUNES

*Departamento de Matemática, Universidade da Beira Interior*  
*Avenida Marquês D'Ávila e Bolama*  
*6201–001 Covilhã, Portugal*  
**e-mail:** celia@noe.ubi.pt

AND

JOÃO TIAGO MEXIA

*Departamento de Matemática, Universidade Nova de Lisboa*  
*Faculdade de Ciências e Tecnologia*  
*Quinta da Torre, 2825–114 Monte da Caparica, Portugal*

### Abstract

Generalized  $F$  tests were introduced by Michalski and Zmysłony (1996) for variance components and later (1999) for linear functions of parameters in mixed linear models. We now use generalized polar coordinates to obtain, for the second case, tests that are more powerful for selected families of alternatives.

**Keywords:** mixed linear models, variance components, generalized polar coordinates, selective  $F$  tests.

**1991 Mathematics Subject Classification:** 62J10, 62J12, 62J99.

### 1. INTRODUCTION

Generalized  $F$  tests were introduced by Michalski and Zmysłony (1996), and (1999) first for variance components and later for linear combinations of parameters in mixed linear models.

We now use polar coordinates to derive generalized  $F$  tests with enhanced power for chosen alternatives. This treatment extends the one we presented for  $F$  tests, see Dias (1994) and Nunes and Mexia (2003).

Our results apply to mixed models satisfying assumptions discussed in the next section. In that section we will also consider the use of polar coordinates in selecting alternatives. In the final section, we discuss the use of polar coordinates in connection with generalized  $F$  statistics.

## 2. MODELS AND HYPOTHESIS

Superscripts will indicate vector dimensions, writing  $Z^n \sim N(\nu^u, W)$  when  $Z^n$  is normal with mean vector  $\nu^u$  and variance-covariance matrix  $W$ . Moreover,  $V \sim \sigma^2 \chi_{g,\delta}^2$  indicates the product by  $\sigma^2$  of a chi-square with  $g$  degrees of freedom and non-centrality parameters  $\delta$ . When  $\delta = 0$  we write simply  $V \sim \sigma^2 \chi_g^2$ .

We will assume that there are mutually orthogonal matrices  $Q_j = A_j^\top A_j$ ,  $j = 1, \dots, J$ , and that, with  $Y^n$  the observations vector, we have the independent vectors such that

$$(1) \quad \begin{cases} A_1 Y^n \sim N(\eta_1^{g_1}, \gamma_1 I_{g_1}) \\ A_j Y^n \sim N(0^{g_j}, \gamma_j I_{g_j}); j = 2, \dots, J \end{cases}$$

where

$$(2) \quad \gamma_1 = \sum_{j=2}^J a_j \gamma_j$$

with  $a_j \leq 0$ ,  $j = 2, \dots, r$ , and  $a_j \geq 0$ ,  $j = r+1, \dots, r+s = J$ .

Thus we have, the independent statistics

$$(3) \quad V_1 = \|A_1 Y^n\|^2 \sim \gamma_1 \chi_{g_1, \delta_1}^2$$

with

$$(4) \quad \delta_1 = \frac{1}{\gamma_1} \|\eta_1^{g_1}\|^2$$

and

$$(5) \quad V_j = \|A_j Y^n\|^2 \sim \gamma_j \chi_{g_j}^2; j = 2, \dots, r+s.$$

Now, the hypothesis

$$(6) \quad H_0 : \eta_1^{g_1} = 0^{g_1}$$

holds if and only if  $\delta_1 = 0$ , and since

$$(7) \quad \begin{cases} E(V_1) = \gamma_1(g_1 + \delta_1) \\ E(V_j) = \gamma_j g_j; \quad j = 2, \dots, r+s \end{cases}$$

if and only if

$$(8) \quad E \left( \frac{V_1}{g_1} - \sum_{j=2}^{r+s} a_j \frac{V_j}{g_j} \right) = 0.$$

We can thus, following Michalski and Zmyslony (1996) and (1999), consider for  $H_0$  a generalized  $F$  test statistic

$$(9) \quad \mathfrak{S} = \frac{\frac{V_1}{g_1} + \sum_{j=2}^r |a_j| \frac{V_j}{g_j}}{\sum_{j=r+1}^{r+s} a_j \frac{V_j}{g_j}},$$

the numerator [denominator] of this statistics being the positive [negative] part of a quadratic unbiased estimator.

As in our previous paper, Nunes and Mexia (2002), polar coordinates  $(r, \Theta^{g_1-1})$  for  $\eta_1^{g_1}$ , will be used to specify selected alternatives for  $H_0$ . Namely we intend to enhance the test power for alternatives such that  $\Theta^{g_1-1} \in \mathcal{C}$  with  $\mathcal{C}$  a conveniently chosen domain. For instance  $\mathcal{C}$  may correspond to certain conditions on the ordering of the components of  $\eta_1^{g_1}$ .

### 3. GENERALIZED $F$ STATISTICS AND POLAR COORDINATES

In defining the selected alternatives to  $H_0$  we will use generalized polar coordinates, see Kendall (1961, p. 15–18). If  $(u, \theta_1, \dots, \theta_{g_1-1})$  are these coordinates we will have, for the components  $\eta_1, \dots, \eta_{g_1}$  of  $\eta_1^{g_1}$

$$(10) \quad \begin{cases} \eta_1 = uc_1 \dots c_{g_1-1} \\ \vdots \\ \eta_i = uc_1 \dots c_{g_1-i} s_{g_1-i+1}; \quad i = 2, \dots, g_1 - 1 \\ \vdots \\ \eta_{g_1} = us_1 \end{cases}$$

with  $u = \|\eta_1^{g_1}\|$ ,  $c_i = \cos\theta_i$  and  $s_i = \sin\theta_i$ ,  $i = 1, \dots, g_1 - 1$ . For the central angles we have the bounds

$$(11) \quad \begin{cases} -\frac{\pi}{2} \leq \theta_i \leq \frac{\pi}{2}; \quad i = 1, \dots, g_1 - 2 \\ 0 \leq \theta_{g_1-1} \leq 2\pi \end{cases}$$

which define the domain  $\mathcal{D}$ .

Selected alternatives will be those for which  $\Theta^{g_1-1} \in \mathcal{C} \subset \mathcal{D}$ , where  $\mathcal{C}$  may be chosen from conditions on the ordering of the  $\eta_1, \dots, \eta_{g_1}$ .

For the components  $Z_1, \dots, Z_{g_1}$  of  $Z^{g_1}$  we have, generalized polar coordinates  $(V_1^{1/2}, \Theta_1, \dots, \Theta_{g_1-1})$ , with

$$(12) \quad \begin{cases} Z_1 = V_1^{1/2} \ell_1(\Theta^{g_1-1}) = UC_1, \dots, C_{g_1-1} \\ \vdots \\ Z_i = V_1^{1/2} \ell_i(\Theta^{g_1-1}) = UC_1 \dots C_{g_1-i} S_{g_1-i+1}; \quad i = 2, \dots, g_1 - 1 \\ \vdots \\ Z_{g_1} = V_1^{1/2} \ell_{g_1}(\Theta^{g_1-1}) = US_1 \end{cases}$$

where  $C_i = \cos\Theta_i$  and  $S_i = \sin\Theta_i$ ,  $i = 1, \dots, g_1$ .

In carrying out selective tests our statistics will be the pair  $(\mathfrak{S}, \Theta^{g_1-1})$ .

To obtain distribution results we start by pointing that, see Kendall (1961, p. 16), the Jacobian of transformations (10) is  $\frac{1}{2}v^{\frac{g_1}{2}-1}h(\theta^{g_1-1})$  with  $h(\theta^{g_1-1}) = c_1^{g_1-2} \dots c_{g_1-2} \geq 0$ .

Since the density of  $Z^{g_1} = A_1 Y^n$  is

$$(13) \quad n(z^{g_1}) = \frac{e^{-\frac{1}{2\gamma_1} \|z^{g_1} - \eta_1^{g_1}\|^2}}{(2\pi)^{g_1/2} \gamma_1^{g_1/2}} = \frac{e^{-\frac{1}{2\gamma_1} (\|z^{g_1}\|^2 - 2\eta_1^\top z + \|\eta_1^{g_1}\|^2)}}{(2\pi)^{g_1/2} \gamma_1^{g_1/2}},$$

when we carry out the transformation for generalized polar coordinates we easily obtain for  $(V_1, \Theta^{g_1-1})$  the joint density

$$\begin{aligned} (14) \quad & f^0(v_1, \theta^{g_1-1} | \delta_1) \\ &= \frac{e^{-\frac{\delta_1}{2}}}{(2\pi)^{g_1/2} \gamma_1^{g_1/2}} h(\theta^{g_1-1}) \frac{v_1^{-1/2}}{2} \sum_{i=0}^{+\infty} \frac{e^{-\frac{v_1}{2\gamma_1}} v_1^{\frac{g_1-1+i}{2}}}{i!} \left( \frac{\eta_1^\top \ell(\theta)}{\gamma_1} \right)^i \\ &= \frac{e^{-\frac{\delta_1}{2}}}{(\pi)^{g_1/2}} \sum_{i=0}^{+\infty} \frac{e^{-\frac{v_1}{2\gamma_1}}}{2\gamma_1 \Gamma(\frac{g_1+i}{2})} \left( \frac{v_1}{2\gamma_1} \right)^{\frac{g_1+i}{2}-1} \frac{\Gamma(\frac{g_1+i}{2}) 2^{i/2-1}}{i! \gamma_1^{i/2}} (\eta_1^\top \ell(\theta))^i h(\theta^{g_1-1}) \\ &= \frac{e^{-\frac{\delta_1}{2}}}{\pi^{g_1/2}} \sum_{i=0}^{+\infty} \frac{1}{\gamma_1} g\left(\frac{v_1}{\gamma_1} | g_1 + i\right) m_i(\theta^{g_1-1}) \end{aligned}$$

where  $m_i(\theta^{g_1-1}) = \frac{\Gamma(\frac{g_1+i}{2}) 2^{i/2-1}}{i! \gamma_1^{i/2}} (\eta_1^\top \ell(\theta^{g_1-1}))^i h(\theta^{g_1-1})$ ,  $i = 0, \dots$  and  $g(x|g_1 + i)$  is the density of a central chi-square with  $g_1 + i$  degrees of freedom. Now, when  $H_0$  holds  $\delta_1 = 0$  as well as  $m_i(\theta^{g_1-1}) = 0$ ,  $i = 1, \dots$ , and so

$$(15) \quad f^0(v_1, \theta^{g_1-1} | 0) = \frac{1}{\pi^{g_1/2}} \frac{1}{\gamma_1} g\left(\frac{v_1}{\gamma_1} | g_1\right) \frac{\Gamma(\frac{g_1}{2})}{2} h(\theta^{g_1-1})$$

thus, when  $H_0$  holds,  $V_1$  and  $\Theta^{g_1-1}$  are independent.

Nextly, since the  $|a_j|V_j$ ,  $j = 2, \dots, r+s$ , have densities  $\frac{1}{|a_j|\gamma_j}g(\frac{v_j}{|a_j|\gamma_j}|g_j)$ ,  $j = 2, \dots, r+s$ , the joint density of  $\Theta^{g_1-1}$ ,  $V_1, \dots, V_{r+s}$  will be

$$\begin{aligned}
 (16) \quad & f^0(v^{r+s}, \theta^{g_1-1}|\delta_1) \\
 &= f^0(v_1, \theta^{g_1-1}|\delta_1) \prod_{i=2}^{r+s} \frac{1}{|a_i|\gamma_i} g\left(\frac{v_i}{|a_i|\gamma_i}|g_i\right) \\
 &= \frac{e^{-\frac{\delta_1}{2}}}{\pi^{g_1/2}} \sum_{i=0}^{+\infty} m_i(\theta^{g_1-1}) \frac{1}{\gamma_1} g\left(\frac{v}{\gamma_1}|g_1+i\right) \prod_{i=2}^{r+s} \frac{1}{|a_i|\gamma_i} g\left(\frac{v_i}{|a_i|\gamma_i}|g_i\right).
 \end{aligned}$$

When  $H_0$  holds we have, with  $a_1 = 1$ ,

$$\begin{aligned}
 (17) \quad & f^0(v^{r+s}, \theta^{g_1-1}|0) \\
 &= \frac{\Gamma(\frac{g_1}{2})}{2\pi^{g_1/2}} \prod_{j=1}^{r+s} \frac{1}{|a_j|\gamma_j} g\left(\frac{v_j}{|a_j|\gamma_j}|g_j\right) h(\theta^{g_1-1}) \\
 &= \left[ \frac{1}{\pi^{g_1/2}} \prod_{j=1}^{r+s} \frac{1}{|a_j|\gamma_j} g\left(\frac{v_j}{|a_j|\gamma_j}|g_j\right) \right] m_0(\theta^{g_1-1}).
 \end{aligned}$$

We thus established

**Proposition 1.** *When  $H_0$  holds  $\mathfrak{S}$  is independent from  $\Theta^{g_1-1}$  which has density  $m_0(\theta^{g_1-1})$ .*

**Proof.** The thesis follows from (17) observing that only  $\mathfrak{S}$  depends on the  $|a_j|V_j$ ,  $j = 1, \dots, r+s$ . ■

Thus, to test  $H_0$  having selected the alternatives for which  $\Theta^{g_1-1} \in \mathcal{C}$  we can reject  $H_0$  whenever  $\Theta^{g_1-1} \in \mathcal{C}$  and  $\mathfrak{S} > k$ . The test level will be the product of  $p(\Theta^{g_1-1} \in \mathcal{C}|H_0)$  by  $p(\mathfrak{S} > k|H_0)$  since, when  $H_0$  holds, both statistics are independent.

When  $\mathcal{C}$  is given by conditions on the ordering of the  $\eta_1, \dots, \eta_{g_1}$  it is usually simple to use these in order to compute  $p(\Theta^{g_1-1} \in \mathcal{C}|H_0)$ , for instance if we require  $\eta_1 = \min\{\eta_1, \dots, \eta_{g_1}\}$ , we have

$$(18) \quad p(\Theta^{g_1-1} \in \mathcal{C}|H_0) = \frac{1}{g_1}.$$

As for the second factor we point out that, when  $H_0$  holds,  $\mathfrak{S}$  is the quotient of two linear combinations  $\sum_{j=1}^r b_j \chi_{g_j}^2$  and  $\sum_{j=r+1}^{r+s} b_j \chi_{g_j}^2$  with  $b_j = \frac{|a_j|}{g_j}$ ,  $j = 1, \dots, r+s$ . Now, Fonseca *et al.* (2002), gave explicit expressions for the distribution of such quotients when all degrees of freedom in the numerator or in the denominator are even. They also showed how to use Monte-Carlo methods in the general case.

For a  $q$  level test we will have

$$(19) \quad q = p(\Theta^{g_1-1} \in \mathcal{C}|H_0)p(\mathfrak{S} > k|H_0)$$

so that, when  $p(\Theta^{g_1-1} \in \mathcal{C}|H_0)$  is small, we are led to choose  $k$  small. Thus a considerable increase of power for selected alternatives is to be expected.

#### REFERENCES

- [1] A. Michalski and R. Zmyślony, *Testing hypothesis for variance components in mixed linear models*, Statistics **27** (1996), 297–310.
- [2] A. Michalski and R. Zmyślony, *Testing hypothesis for linear functions of parameters in mixed linear models*, Tatra Mountain Mathematical Publications **17** (1999), 103–110.
- [3] C. Nunes and J.T. Mexia, *Selective  $F$  tests for sub-normal models*, Discussiones Mathematicae, Probability and Statistics **23** (2003), 167–174.
- [4] G. Dias, *Selective  $F$  tests*, Trabalhos de Investigação N<sup>o</sup> 1, FCT/UNL (1994).

- [5] M. Fonseca, J.T. Mexia and R. Zmysłony, *Exact distribution for the generalized  $F$  tests*, *Discussiones Mathematicae, Probability and Statistics* (1,2) **22** (2002), 37–51.
- [6] M.G. Kendall, *A Course in the Geometry of  $n$  Dimensions*, Charles Griffin & Company Limited, London 1961.

Received 1 June 2004