

Discussiones Mathematicae
Probability and Statistics 34 (2014) 63–69
doi:10.7151/dmps.1170

ON USEFUL SCHEMA IN SURVIVAL ANALYSIS AFTER HEART ATTACK

CZESŁAW STĘPNIAK

Department of Differential Equations and Statistics
Faculty of Mathematics and Natural Sciences
University of Rzeszów
Pigonia 1, 35–959 Rzeszów, Poland
e-mail: stepniak@umcs.lublin.pl

Abstract

Recent model of lifetime after a heart attack involves some integer coefficients. Our goal is to get these coefficients in simple way and transparent form. To this aim we construct a schema according to a rule which combines the ideas used in the Pascal triangle and the generalized Fibonacci and Lucas numbers

Keywords: lifetime after heart attack, distribution, Fibonacci number, Lucas number, Pascal triangle.

2010 Mathematics Subject Classification: Primary 62E15, 60C05; Secondary 05A10, 11B39.

1. INTRODUCTION

This note is motivated by a real medical problem. Many doctors believe that a patient survives a heart attack (in medical terminology: myocardial infarction) unless a succeeding attack occurs in a week. The length of this critical period (for definition see [12] p. 372) is still controversial. Treating heart attacks as failures in Bernoulli trials we reduce the lifetime after a heart attack to the waiting time for the first failure followed by a success run shorter than a given k . Its distribution, recently presented in [25], involves some integer coefficients. Our goal is to get these coefficients both in a simple way and in a transparent form. For this aim

we construct a schema according to a rule which combines the ideas used in the Pascal triangle and the generalized Fibonacci and Lucas numbers.

Systems of positive integers have been investigated very intensively in the last decade. In fact this area become a specialized branch of discrete mathematics represented, among others, by five books [10, 19, 27, 28] and [15], two specialized journals (Fibonacci Quarterly and Journal of Integer Sequences) and the On-Line Encyclopedia of Integer Sequences [22]. The main attention focused on Fibonacci, Lucas and Catalan Numbers (FNs, LNs and CNs for abbreviation) with possible generalizations (cf. [3, 4, 5, 9, 14, 17, 20, 24, 30] and [7]) on polynomials generated by them (cf. [2, 3, 6, 18, 23, 26] and [1]). A nice geometric interpretation both FNs and LNs within the Pascal triangle (PT) has been revealed by Koshy [16]. In recent works [13] and [29] FNs, LNs and CNs were studied in terms of graph theory.

This note is the next in the series of works [8, 11, 21] and [13] on schemata of positive integers. It is a response to the real need.

2. FROM DISTRIBUTION OF LIFETIME AFTER HEART ATTACK TO SCHEMA OF ITS COEFFICIENTS

The waiting time for the first failure followed by a success run shorter than k may be expressed by the following model.

Let $X = (X_1, X_2, X_3, \dots)$ be a sequence of independent identically distributed random variables taking values 1 or 0 with probabilities p and $q = 1 - p$, where 0 is interpreted as a failure. Given a positive integer k called critical period, define a statistic $T = t(X)$ as the minimal integer n such that $X_n = 0$ and, either $n \leq k$ or $X_m = 0$ for some m satisfying the condition $0 < n - m \leq k$. The statistic T is said to be waiting time for the first failure followed by a success run shorter than k .

Recently the probability mass function of the waiting time in this model was derived by Stępnia [25] in the form

$$P(T = n) = \begin{cases} p^n t, & \text{if } n = 1, 2, \dots, k, \\ (n - k - 1)p^n t^2, & \text{if } n = k + 1, \dots, 2k + 1, \\ p^n t^2 \sum_{r=0}^{\left[\frac{n-k-2}{k+1}\right]} a_r t^r, & \text{if } n > 2k + 1, \end{cases}$$

where $t = \frac{q}{p}$, $[x]$ means the integer part of a number x , while a_r , for $r = 0, \dots, \left[\frac{n-k-2}{k+1}\right]$, is the system of the integer coefficients defined by

$$(1) \quad a_r = a_{n,r;k} = \binom{n - (r+1)k - 1}{r+1} - \binom{n - (r+2)k - 1}{r+1}.$$

It is well known (cf. e.g. [7, 18, 20, 26]) that the generalized FNs and LNs operate with the same recurrence relation

$$(2) \quad N(n) = N(n-1) + N(n-k-1),$$

but with different initial conditions. On the other hand the construction of the Pascal triangle with its entries

$$(3) \quad \binom{n}{r} = \begin{cases} \frac{n!}{r!(n-r)!}, & \text{if } 0 \leq r \leq n \\ 0, & \text{otherwise} \end{cases},$$

is based on the relation

$$(4) \quad \binom{n}{r} = \binom{n-1}{r-1} + \binom{n-1}{r}$$

with the initial conditions

$$\binom{n}{r} = \begin{cases} 1, & \text{if } r = 0 \\ 0, & \text{if } n = 1 \text{ and } r > 0. \end{cases}$$

This construction may be explained by the following schema:

		0		r-1		r			
0		1							...
		1	1						...
		1	2	1					...
n-1		1	3	3	1				...
n		1	4	6	4	1			...
		1	5	10	10	5	1		...
		1	6	15	20	15	6	1	...
	...	...	...	...	...	...	...	...	...

Combining (2) and (4) one can get a k -step recurrence relation of type

$$\binom{n}{r}_k = \binom{n-1}{r}_k + \binom{n-k-1}{r-1}_k.$$

This relation may be also completed by some initial condition (as in the generalized FNs and LNs).

It appears that a similar rule may be used to the system of the coefficients in distribution of the lifetime after heart attack.

Definition 1. The system of the coefficients $a_{n,r;k}$ defined by the formula (1) for a given integer $k \geq 0$ and for $n, r = 1, 2, \dots$ is said to be k -schema.

Remark 2. For $k = 0$ the k -schema coincides with the usual Pascal triangle.

Theorem 3. The entries $a_{n,r;k}$ in k -schema satisfy the recurrence relation

$$(5) \quad a_{n,r;k} = a_{n-1,r;k} + a_{n-k-1,r-1;k}$$

for $n \geq k + 2$ and $r \geq 1$ with the initial conditions

$$a_{n,r;k} = \begin{cases} 0, & \text{if } n < k + 2 \\ n - k - 1, & \text{if } k + 2 \leq n < 2k + 2 \text{ and } r = 0 \\ k, & \text{if } n \geq 2k + 2 \text{ and } r = 0. \end{cases}$$

Proof. Let us start from the initial conditions.

By (1) and (3), $a_{r,n;k} = 0$ if and only if $\binom{n-(r+1)k-1}{r+1} = 0$, i.e., when $n - 1 < (r + 1)(k + 1)$; in particular when $n < k + 2$.

If $n \in \{k + 2, \dots, 2k + 1\}$ then $n < 2k + 2$. In consequence

$$a_{n,0;k} = \binom{n - k - 1}{1} - \binom{n - 2k - 1}{1} = n - k - 1.$$

Now let $n \geq 2k + 2$. Then

$$a_{n,0;k} = \binom{n - k - 1}{1} - \binom{n - 2k - 1}{1} = n - k - 1 - (n - 2k - 1) = k.$$

For relation (5) we shall use the property (4). By definition (1)

$$\begin{aligned} a_{n-1,r;k} &= \binom{n - 1 - (r + 1)k - 1}{r + 1} - \binom{n - 1 - (r + 2)k - 1}{r + 1} \\ &= \binom{n - (r + 1)k - 2}{r + 1} - \binom{n - (r + 2)k - 2}{r + 1} \end{aligned}$$

and

$$\begin{aligned} a_{n-k-1,r-1;k} &= \binom{n-k-1-rk-1}{r} - \binom{n-k-1-(r+1)k-1}{r} \\ &= \binom{n-(r+1)k-2}{r} - \binom{n-(r+2)k-2}{r}. \end{aligned}$$

Thus by (1) and (4) we get (5). ■

Theorem 3 constitutes a basis of our schema. For example, if $k = 3$, this schema takes the following form

	0	r-1	r	...
0	0			...
	0			...
	0			...
	0			...
	0			...
	1			...
	2			...
	3			...
	3			...
	3	1		...
	3	3		...
n-k-1	3	6		...
	3	9		...
	3	12	1	...
n-1	3	15	4	...
n	3	18	10	...
...	...	...	...	...

Acknowledgement

This work was partially supported by the Centre for Innovation and Transfer of Natural Sciences and Engineering Knowledge.

REFERENCES

- [1] H. Belbachir and A. Benmezai, *An alternative approach to Cigler's q -Lucas polynomials*, Appl. Math. Computat. **226** (2014) 691–698. doi:10.1016/j.amc.2013.10.009
- [2] G.B. Diordjević, *Generating functions of the incomplete generalized Fibonacci and generalized Lucas numbers*, Fibonacci Quart. **39** (2004) 106–113.

- [3] A. Dil and I. Mező, *A symmetric algorithm for hyperharmonic and Fibonacci numbers*, Appl. Math. Comput. **206** (2008) 942–951. doi:10.1016/j.amc.2008.10.013
- [4] M. El-Mikkawy and T. Sogabe, *A new family of k -Fibonacci numbers*, Appl. Math. Comput. **215** (2010) 4456–4461. doi:10.1016/j.amc.2009.12.069
- [5] X. Fu and X. Zhou, *On matrices related with Fibonacci and Lucas numbers*, Appl. Math. Comput. **200** (2008) 96–100. doi:10.1016/j.amc.2007.10.060
- [6] D. Garth, D. Mills and P. Mitchell, *Polynomials generated by the Fibonacci sequence*, J. Integer. Seq. **10** (2007), Article 07.6.8.
- [7] H.H. Gulec, N. Taskara and K. Uslu, *A new approach to generalized Fibonacci and Lucas numbers with binomial coefficients*, Appl. Math. Comput. **230** (2013) 482–486. doi:10.1016/j.amc.2013.05.043
- [8] J.M. Gutiérrez, M.A. Hernández, P.J. Miana and N. Romero, *New identities in the Catalan triangle*, J. Math. Anal. Appl. **341** (2008) 52–61. doi:10.1016/j.jmaa.2007.09.073
- [9] P. Hao and S. Zhi-wei, *A combinatorial identity with application to Catalan numbers*, Discrete Math. **306** (2006) 1921–1940. doi:10.1016/j.disc.2006.03.050
- [10] V.E. Hoggat Jr., *Fibonacci and Lucas Numbers*, Houghton Mifflin (Boston, MA, 1969).
- [11] H. Hosoya, *Fibonacci triangle*, Fibonacci Quart. **14** (1976) 173–178.
- [12] B.D. Jones, *Comprehensive Medical Terminology*, Third Ed. Delmar Publishers (Albany NY, 2008).
- [13] S. Kitaev and J. Liese, *Harmonic numbers, Catalan's triangle and mesh patterns*, Discrete Math. **313** (2013) 1515–1531. doi:10.1016/j.disc.2013.03.017
- [14] E.G. Kocer and N. Touglu, *The Binet formulas for the Pell-Lucas p -numbers*, Ars Combinatoria **85** (2007) 3–18.
- [15] T. Koshy, *Fibonacci and Lucas Numbers with Applications* (Wiley-Interscience, New York, 2001). doi:10.1002/9781118033067
- [16] T. Koshy, *Fibonacci, Lucas, and Pell numbers, and Pascal's triangle*, Math. Spectrum **43** (2011) 125–132.
- [17] H. Kwong, *Two determinants with Fibonacci and Lucas entries*, Appl. Math. Comput. **194** (2007) 568–571. doi:10.1016/j.amc.2007.04.027
- [18] S.-M. Ma, *Identities involving generalized Fibonacci-type polynomials*, Appl. Math. Comput. **217** (2011) 9297–9301. doi:10.1016/j.amc.2011.04.012
- [19] L. Niven, H. Zuckerman and H. Montgomery, *An Introduction to the Theory of Numbers*, Fifth Ed. (Wiley, New York, 1991).
- [20] J. Petronilho, *Generalized Fibonacci sequences via orthogonal polynomials*, Appl. Mat. Comput. **218** (2012) 9819–9824. doi:10.1016/j.amc.2012.03.053

- [21] L.W. Shapiro, *A Catalan triangle*, Discrete Math. **14** (1976) 83–90.
doi:10.1016/0012-365X(76)90009-1
- [22] N. Sloane, On-Line Encyclopedia of Integer Sequences (OEIS), <http://oeis.org>.
- [23] S. Stanimirović, *Some identities on Catalan numbers and hypergeometric functions via Catalan matrix power*, Appl. Math. Comput. **217** (2011) 9122–9132.
doi:10.1016/j.amc.2011.03.138
- [24] S. Stanimirović, P. Stanimirović, M. Miladinović and A. Ilić, *Catalan matrix and related combinatorial identities*, Appl. Math. Comput. **215** (2009) 796–805.
doi:10.1016/j.amc.2009.06.003
- [25] C. Stepniak, *On distribution of waiting time for the first failure followed by a limited length success run*, Appl. Math. (Warsaw) (2013) 421–430. doi:10.4064/am40-4-3
- [26] N. Tuglu, E.G. Kocer and A. Stakhov, *Bivariate fibonacci like p -polynomials*, Appl. Math. Comput. **217** (2011) 10239–10246. doi:10.1016/j.amc.2011.05.022
- [27] S. Vajda, *Fibonacci and Lucas Numbers and the Golden Section*, Ellis Horwood (Chichester 1989).
- [28] N.N. Vorobyov, *Fibonacci Numbers*, Publishing House "Nauka", Moscow, 1961 (in Russian).
- [29] A. Włoch, *Some identities for the generalized Fibonacci numbers and the generalized Lucas numbers*, Appl. Math. Comput. **219** (2013) 5564–5568.
doi:10.1016/j.amc.2012.11.030
- [30] O. Yayenie, *A note on generalized Fibonacci sequences*, Appl. Math. Comput. **217** (2011) 5603–5611. doi:10.1016/j.amc.2010.12.038

Received 19 June 2014

