

*Discussiones Mathematicae*  
*Probability and Statistics* 30 (2010) 117–122  
doi:10.7151/dmps.1124

## SAMPLE PARTITIONING ESTIMATION FOR ERGODIC DIFFUSIONS: APPLICATION TO ORNSTEIN-UHLENBECK DIFFUSION

LUÍS RAMOS

*Department of Mathematics*  
*New University of Lisbon, Portugal*

**e-mail:** lpcr@fct.unl.pt

### Abstract

When a diffusion is ergodic its transition density converges to its invariant density, see Durrett (1998). This convergence enabled us to introduce a sample partitioning technique that gives in each sub-sample, maximum likelihood estimators. The averages of these being a natural choice as estimators. To compare our estimators with the optimal we obtained from martingale estimating functions, see Sørensen (1998), we used the Ornstein-Uhlenbeck process for which exact simulations can be carried out.

**Keywords:** ergodic diffusions; martingale estimating functions; transition and invariant densities; maximum likelihood estimators.

**2000 Mathematics Subject Classification:** 60J60, 60H35, 62F10.

### REFERENCES

- [1] B.M. Bibby and M. Sørensen, *Martingale estimation functions for discretely observed diffusion processes*, Bernoulli **1** (1995), 17–39. doi:10.2307/3318679
- [2] U. Küchler and M. Sørensen, *Exponential Families of Stochastic Processes*, Springer-Verlag 1997.
- [3] S. Iacus, *Simulation and Inference for Stochastic Differential, Equations with R Examples*, Springer 2008. doi:10.1007/978-0-387-75839-8

- [4] J.T. Mexia and G.C. Dias, Statistical Inference for Discretely Observed Diffusions of Diffusions, VII-Congresso da SPE, 14–16 Outubro, Osir 1999.
- [5] B. Øksendal, *Stochastic Differential Equations*, An Introduction, Fifth Edition, Springer-Verlag 1998.
- [6] M. Sørensen, Lecture Notes on "Statistical Inference for Discretely Observed Diffusions", Berlin Graduiertenkolleg 1998.

Received 18 January 2010